

IN THE CLAIMS:

Please amend the claims to provide as follows.

Claims 1-10 (Cancelled).

Claim 11. (Previously Added) A method for transduction of hematopoietic cells by a replication-defective recombinant retrovirus vector, comprising infecting viable hematopoietic cells in culture with a replication-defective recombinant retrovirus in the presence of substantially pure fibronectin, substantially pure fibronectin fragments, or a mixture thereof, to produce transduced hematopoietic cells, said fibronectin and said fibronectin fragments containing the alternately spliced CS-1 cell adhesion domain and the Heparin II binding domain of fibronectin.

Claim 12. (Previously Added) The method of claim 11 which includes harvesting the transduced hematopoietic cells.

Claim 13. (Previously Added) The method of claim 11 wherein the hematopoietic cells have a protein deficiency or abnormality and the recombinant retrovirus vector includes an exogenous gene encoding the protein.

Claim 14. (Previously Added) The method of claim 11 wherein the hematopoietic cells have an enzyme deficiency or abnormality and the exogenous gene is a gene encoding the enzyme.

Claim 15. (Previously Added) The method of claim 14 wherein the hematopoietic cells are human hematopoietic cells having an enzyme deficiency or abnormality and the exogenous gene is a human gene encoding the enzyme.

Claim 16. (Previously Added) The method of claim 14 wherein the hematopoietic cells have an adenosine deaminase deficiency and the exogenous gene encodes adenosine deaminase.

Claim 17. (Previously Added) The method of claim 15 wherein the human hematopoietic cells have an adenosine deaminase deficiency and the exogenous gene encodes adenosine deaminase.

Claim 18. (Previously Added) The method of claim 15 wherein the cells are infected with the retrovirus in the presence of an immobilized fibronectin fragment containing an amino acid sequence which provides the cell-binding activity of the CS-1 domain and an amino acid sequence which provides the retrovirus binding activity of the Heparin-II domain.

Claim 19. (Previously Added) The method of claim 18 wherein the fibronectin fragment is a recombinant fibronectin fragment.

Claim 20. (Previously Added) The method of claim 19, wherein the recombinant fibronectin fragment is selected from the group consisting of H-296 and CH-296.

Claim 21. (Previously Added) The method of claim 20, wherein the recombinant fibronectin fragment is CH-296.

Claim 22. (Previously Added) The method of claim 19, wherein the recombinant fibronectin fragment contains the Heparin-II binding domain of fibronectin.

Claim 23. (Previously Added) The method of claim 11, wherein the hematopoietic cells are characterized as adherent-negative, low density, mononuclear cells.

Claims 24-37 (Cancelled).

Claim 38. (Currently Amended) A cellular composition comprising viable hematopoietic cells transduced by retroviral-mediated gene transfer in the absence of

retroviral producer cells and in the presence of an immobilized amount of a polypeptide containing fibronectin, a fibronectin fragment, or a mixture thereof, a first amino acid sequence which provides the binding activity of the Heparin-II binding domain of fibronectin and a second amino acid sequence which provides the cell-binding activity of the CS-1 domain of fibronectin, said immobilized amount of polypeptide being effective to increase the frequency of transduction of the hematopoietic cells by the retrovirus vector; said composition also comprising said polypeptide.

Claim 39. (Previously Added) The cellular population of claim 38 which is enriched in hematopoietic stem cells.

Claim 40. (Previously Added) The cellular population of claim 38 wherein said viable hematopoietic cells are human hematopoietic cells enriched in human hematopoietic stem cells.

Claim 41. (Previously Added) The cellular population of claim 40 which is a substantially homogenous population of human hematopoietic cells characterized as adherent-negative, low density, mononuclear cells.

Claim 42. (Previously Added) The cellular population of claim 41 which has been transduced by a recombinant retrovirus vector containing an exogenous gene to correct a protein deficiency or abnormality in the cells.

Claim 43. (Previously Added) The cellular population of claim 39 wherein said hematopoietic cells are obtained from umbilical cord blood.

Claims 44-78 (Cancelled).

Claim 79. (Previously Added) In a method of gene transfer into mammalian cells by a replication-defective recombinant retrovirus vector, the improvement comprising conducting the gene transfer without cocultivation and in the presence of substantially pure fibronectin, substantially pure fibronectin fragments, or a mixture thereof, so as to increase the frequency of the gene transfer.

Claim 80. (Previously Added) A method for transduction of viable mammalian cells by a replication-defective recombinant retrovirus vector, comprising infecting the cells in culture with a replication-defective recombinant retrovirus in the presence of substantially pure fibronectin, substantially pure fibronectin fragments, or a mixture thereof, to produce transduced cells.

Claim 81. (Previously Added) The method of claim 80, wherein the infecting is in the presence of a fibronectin fragments containing the Heparin-II binding domain of fibronectin.

Claim 82. (Previously Amended) The method of claim 81, wherein said domain has an amino acid sequence represented by the formula (SEQ. ID NO. 1):

Ala Ile Pro Ala Pro Thr Asp Leu Lys Phe Thr Gln Val Thr Pro Thr Ser Leu Ser Ala  
Gln Trp Thr Pro Pro Asn Val Gln Leu Thr Gly Tyr Arg Val Arg Val Thr Pro Lys Glu  
Lys Thr Gly Pro Met Lys Glu Ile Asn Leu Ala Pro Asp Ser Ser Ser Val Val Val Ser  
Gly Leu Met Val Ala Thr Lys Tyr Glu Val Ser Val Tyr Ala Leu Lys Asp Thr Leu  
Thr Ser Arg Pro Ala Gln Gly Val Val Thr Thr Leu Glu Asn Val Ser Pro Pro Arg  
Arg Ala Arg Val Thr Asp Ala Thr Glu Thr Thr Ile Thr Ile Ser Trp Arg Thr Lys Thr  
Glu Thr Ile Thr Gly Phe Gln Val Asp Ala Val Pro Ala Asn Gly Gln Thr Pro Ile Gln  
Arg Thr Ile Lys Pro Asp Val Arg Ser Tyr Thr Ile Thr Gly Leu Gln Pro Gly Thr Asp  
Tyr Lys Ile Tyr Leu Tyr Thr Leu Asn Asp Asn Ala Arg Ser Ser Pro Val Val Ile Asp  
Ala Ser Thr Ala Ile Asp Ala Pro Ser Asn Leu Arg Phe Leu Ala Thr Thr Pro Asn  
Ser Leu Leu Val Ser Trp Gln Pro Pro Arg Ala Arg Ile Thr Gly Tyr Ile Ile Lys Tyr  
Glu Lys Pro Gly Ser Pro Pro Arg Glu Val Val Pro Arg Pro Arg Pro Gly Val Thr Glu  
Ala Thr Ile Thr Gly Leu Glu Pro Gly Thr Glu Tyr Thr Ile Tyr Val Ile Ala Leu Lys  
Asn Asn Gln Lys Ser Glu Pro Leu Ile Gly Arg Lys Lys Thr.

Claim 83. (Currently Amended) The method of claim 82, wherein said fibronectin fragments comprise a recombinant fibronectin fragment selected from the group consisting of CH-296 and H-296.

Claims 84-93 (Cancelled).